



## PHYSICAL PROPERTIES AND SENSORY EVALUATION OF ICE CREAM FORTIFIED WITH CICHORIUM PUMILUM, JACQ CRUDE EXTRACT

**Dr. Nadia A. Abou-zeid\***

Dairy Science and Technology Department, Faculty of Agriculture, Menoufia University,  
Shebin El-kom, Egypt.

Article Received on 01/12/2019

Article Revised on 22/12/2019

Article Accepted on 12/01/2020

### \*Corresponding Author

**Dr. Nadia A. Abou-Zeid**

Dairy Science and  
Technology Department,  
Faculty of Agriculture,  
Menoufia University,  
Shebin El-Kom, Egypt.

### ABSTRACT

The genus Cichorium has a rich history of use in folklore and is widely eaten by the farmers. It is also widely used medicinally to treat various ailments ranging from wounds to diabetes. It was discovered that the root of Cichorium contained up to 40% inulin, which has a negligible impact on blood sugar and thus is suitable for diabetics, the crude extract prepared from cichorium had satisfactory proteolytic activities

which offers a practical way to reduce its antigenic protein fractions.

In this study we investigated the effects of fortified milk (that is used in ice cream manufacture) with Cichorium crude extract (CCA) on the physical and sensory properties of the resulting ice cream.

### INTRODUCTION

Dietary factors are implicated in the etiology of a number of chronic degenerative diseases (Harrigan, 1989; Haumann, 1986). Health-conscious consumers continue to look for ways to improve nutritional habits without sacrificing psychological satisfaction (Kostias, 1997; O'Brien, 2003; Plug, 1993).

Ice cream is a frozen foam that consists of air cells dispersed in an aqueous matrix (Marshall et al., 2003). The three main structural components of ice cream are air cells, ice crystals, and fat globules, which are distributed throughout a continuous phase of unfrozen solution

(serum). Overall, consumers are quite concerned with the influence of diet on health. Eating ice cream has great benefits such as increasing insulin secretion and increasing the gut hormone glucagon-like peptide-1 (GLP-1) which is an important regulator of glucose homeostasis and energy balance through its concerted actions on appetite, Hans (2008), but ice cream, have been considered to increase the risk for allergy since beta-lactoglobulin can induce milk allergy (Kim et al 2007). Enzymatic hydrolysis of milk whey protein offers a practical way to reduce its antigenic protein fractions (Kim et al 2007 and Hai et al 2008), Our previous study found that Cichorium crude extract (CCPE) has proteolytic activity, Nadia (2015).

Cichorium intybus is a medicinally important plant in Eurasia and in parts of Africa. Several studies have been conducted on grazing animals to determine the antihelmintic potential of Cichorium. In the majority of the experiments, the condensed tannins and sesquiterpene lactones were responsible for the anthelmintic activity [D.Mares et al 2011]. It was discovered that the root of Cichorium contained up to 40% inulin, which has a negligible impact on blood sugar and thus is suitable for diabetics [A. Judzientienė and J.B. Audienė 2008], The methanolic extract displays wound healing effect and  $\beta$ -sitosterol was determined as the active compound responsible for the activity, possibly due to its significance. The crude ethanolic extract of C. roots caused a significant inhibition of Ehrlich tumor carcinoma in mice. A 70% increase in the life span was observed with a 500mg/kg/day intraperitoneal dose of the tested extract [B. Hazra et al 2002]. Cichorium crude extracts have anti-inflammatory and antioxidant effects, as well as hyaluronidase and collagenase inhibition [I. Šuntar et al 2012]. Cichorium crude aqueous and organic seed extracts were found to be active against four pathogenic microorganisms, namely: Staphylococcus aureus, Pseudomonas aeruginosa, Escherichia coli, and Candida albicans, and root extracts had pronounced effects on Bacillus subtilis, S. aureus, Salmonella typhi, Micrococcus luteus, and E. coli [I. Šuntar et al 2012, S. Nandagopal 2007]. The flowers of chicory contain saccharides, methoxycoumarin cichorine, flavonoids, essential oils [A. Judzientienė and J.B. Audienė 2008]. The leaf extract of Cichorium also showed a moderate activity against multidrug resistant S. typhi [P. Rani et al 2004]. Guaianolides-rich root extracts of C. intybus have shown antifungal properties against anthropophilic fungi Trichophyton tonsurans, T. rubrum, and T. violaceum [D.Mares et al 2011]. Extraction of chicory pulp yields high levels of pectin, a polysaccharide extensively used in food as a gelling agent, thickening agent, and stabilizer [C. Robert et al 2008].

The crude protein content in chicory is more valuable than in alfalfa. Furthermore the crude lipid in chicory is generally higher than most varieties of alfalfa [Q. Wang and J. Cui, 2011]. It also has a nutritional quality comparable to lucerne, with a mineral content superior for copper and zinc, with similar proportions of protein, lipid, minerals, and other nutrients, and palatable with good digestion with applicability in the poultry and livestock industry [M. A. Sanderson et al 2003, A. Scharenberg et al 2007].

Cichorium is safe for human use, The toxicity evaluation of Cichorium extracts has also been done by *Vibrio fischeri* bioluminescence inhibition test (Microtox acute toxicity test). This bacterial test measures the decrease in light emission from the marine luminescent bacteria *V. fischeri* when exposed to organic extracts. The tested extracts showed less than 20% inhibition of bioluminescence and hence were concluded to be safe for human use [F. Conforti 2008].

Fortification of the milk base is one of the most important steps that enhances functional and nutritional Properties Marie et al (2013) This research utilizes these approaches to produce ice cream fortified with cichorium crude extract to produce functional ice cream with many health benefits.

## **MATERIAL AND METHODS**

### **Preparation of Cichorium crude extract**

Dried whole plant was cut into small pieces grind , maceration 10 gm of this powder in 50 ml hot water (60 c) using an electric mixer . The extract filtered and centrifuged at 30000 rpm for 30min. The supernatant was referred as cichorium crude extract (CCA).

### **Processing**

Vanilla ice cream mixes were made in 3-kg batches as described by Nadia( 2008 ) Fresh bulk buffalo's milk obtained from the farm of the Agriculture faculty ,Minufiyya Univ. ,Shibin El Kom, Egypt.

The ice cream formulations contained 12% milk fat, 16% sucrose, 0.5% stabilizer.

Cichorium crude extract added to the pasteurized milk that will be used in ice cream manufacture at a levels of 2,4,6,8, and 10% batches named CCA2; CCA4 ; CCA6; CCA8 and CCA10 respectively and kept at room temperature for four hours , then used for ice cream mix preparation. All mixes pasteurized for 30 mins. at 71°C .The mixes were then cooled and

aged at 4 to 5°C for 12 h. The ice cream was frozen in a batch freezer (Cattabriga, Bologna, Italy) samples were tempered in a freezer at -15° for approximately 18 h.

### Ice cream analysis

#### Overrun

One overrun measurement was taken per sample by comparing the weight of ice cream mix and ice cream in a fixed volume container. Overrun (in %) was calculated as follows. A plastic whipping bowl was modified to test foam stability (Lim et al., 2008). The time to attain 50% drainage was used as an index of foam stability (Phillips et al., 1990).

$$\text{Overrun} = \frac{\text{weight of mix} - \text{weight of ice cream}(100)}{\text{weight of ice cream}}$$

#### Melting rate

A sample of ice cream was cut from the end of a rectangular half-gallon container to make a rectangular block of ice cream that weighed  $100 \pm 1$  g. The dimensions of the sample varied slightly from one ice cream to another (6 to 7 cm x 6 to 7 cm x 3 cm) due to the differences in physical characteristics (e.g., overrun). The sample of ice cream (initially at -15°C) was placed on a wire screen (6 holes/cm) on top of a funnel that was attached to a graduated cylinder. The ice cream was placed in a controlled temperature chamber at 25°C (RH 50%). Every 5 min, for up to 3 h, the dripped volume was recorded. Pictures of the ice cream were taken with a digital camera every 20 min. The time (min) was plotted against the dripped volume (mL) and the slope of the main melting event was taken as the melting rate (Muse M. R. and. Hartel R. W.,(2004).

#### Hardness

Texture analyzer CNS – (The Farnell, England) was used to measure the hardness of the ice cream at -15°C. Samples were quickly transported to the texture analyzer, held at room temperature, and analyses completed within 30 s to minimize variability due to sample warming. A 42-g stainless steel probe (60° cone with a maximum diameter of 6.5 cm) was aligned so that it touched the surface of the ice cream and then allowed to penetrate the samples under the force for five second. The forces needed for make equal penetration was directly proportional to the hardness. Three measurements were taken per sample. (Muse M. R. and. Hartel R. W, (2004).

### Sensory evaluation

Sensory evaluation consisting of flavor & taste, body&texture, and overall acceptability was carried out by 15 panelists from dairy and food sciences departments.

### Statistical analysis

All the obtained data were statistically analyzed by SPSS computer software according to (Abo- Allam, 2003).

## RESULTS AND DISCUSSION

No significant difference was observed between melting resistance of samples of the different level of CCA until 4% and the control ( $P > 0.05$ ) but the melting resistance of samples containing 6, 8 and 10% significantly increased (Table 1).

As can be seen from Table 1, using CCA at level 6, 8 or 10 % led to increasing hardness of the product. In this case control samples showed lower values of hardness than those with CCA. Increasing CCA level to 4% had no significant effect on hardness of samples.

The increase of instrumental hardness with increased CCA content may be attributed to the freeze concentration in their serum phase (Bolliger et al., 2000).

Viscosity values (Table 2) increased with an increase in the level of CCA but the differences were not significant until 4%. The function of CCA in ice cream preparation is considered to be related to its water binding capacity by forming a three-dimensional network of hydrated molecules throughout the system. Water retention is associated with the mobility of the free water molecules to become limited and consequently the bulk viscosity increased. In this way they retard ice crystal formation and growth, improving mix viscosity, air incorporation, body and texture as well as melting properties (Nielsen, 1978). Overrun (Table 2) significantly increased in samples contain CCA at a level above 4% with no significant differences between control and samples containing 2 and 4 %.

Sensory properties of ice cream containing CCA at five levels are presented in (Table 3). As seen from the table, using CCA at any levels until 6% did not have any negative effect on these properties and furthermore the flavor, texture, color and overall acceptability of ice cream samples containing CCA, were more appealing than control. Considering the scores of all sensory properties, the most acceptable sample was ice cream containing 6% CCA. Texture, Flavor, color and overall acceptance of ice cream containing the two levels 4 and 6%

of CCA did not significantly differ ( $P > 0.05$ ). Taste & flavor scores of the samples containing 8 and 10% were somehow lower than for the other samples and control. They had bitter taste.

**Table (1): Melting rate, hardness in ice creams.**

<b>Determinations Samples*</b>	<b>Melting rate (ml/min)</b>	<b>Hardness(gm) (Depth of penetration 7mm)</b>
C	1.9	131
CCA2	1.9	131
CCA4	1.8	133
CCA6	1.5	146
CCA8	1.2	157
CCA10	1.0	164

\*C: control; CCA2; CCA4; CCA6; CCA8 and CCA10: ice cream fortified with 2,4,6,8, and 10% Cichorium crude extract.

**Table (2): Apparent viscosity overrun in ice cream.**

<b>Determinations Samples*</b>	<b>Apparent Viscosity at 20 rpm (cP)</b>	<b>Overrun %</b>
C	886	75.6
CCA2	889	76.5
CCA4	898	78.1
CCA6	912	93.1
CCA8	958	95.9
CCA10	989	98.4

\*for details see table 1

**Table (3): Organoleptic properties of ice cream.**

<b>Ice cream</b>	<b>Flavour&amp;taste (40)</b>	<b>Body and texture (30)</b>	<b>overall acceptability (30)</b>	<b>Total score (100)</b>
C	40	28	28	96
CCA 2	40	29	27	96
CCA4	40	28	29	97
CCA6	38	30	29	97
CCA8	32	29	26	87
CCA10	30	28	25	73

\*for details see table 1

## REFERENCES

1. Abo-Allam, R. M. Data Statistical Analysis Using SPSS Program. 1st Ed., Publication for Universities, Cairo, 2003.

2. Bolliger, S., Wildmoser, H., Goff, H. D. and Tharp, B. W. Relationships between Ice Cream Mix Viscosity and Ice Crystal Growth in Ice Cream. *Int. Dairy J.*, 2000; 10: 791–797.
3. F. Conforti, G. Ioele, G. A. Statti, M. Marrelli, G. Ragno, and F. Menichini. “Antiproliferative activity against human tumor cell lines and toxicity test on Mediterranean dietary plants,” *Food and Chemical Toxicology*, 2008; 46(10): 3325–3332.
4. Hai Zheng, Xiaoqin Shen, Guanhao Bu and Yongkang Luo Effects of pH, temperature and enzyme to substrate ratio on antigenicity of whey protein hydrolysates prepared by alcalase, *International Dairy Journal*, 2008; 18(10-11): 1028-1033.
5. Hans-Rudolf Berthoud Paying the Price for Eating Ice Cream: Is Excessive GLP-1 Signaling in the Brain the Culprit? *Endocrinology*, 2008; 149(10): 4765-7.
6. Harrigan, K.A., Breene, W.M. Fat substitutes: Sucrose esters and simplese, *Cereal Foods World*, 1989; 34: 261-267.
7. Haumann, B.F. Getting the fat out – researchers seek substitutes for full-fat fat, *Journal of American Oil Chemistry Society*, 1986; 63: 278-286.
8. Hazra, R. Sarkar, S. Bhattacharyya, and P. Roy, “Tumour inhibitory activity of chicory root extract against Ehrlich ascites carcinoma in mice,” *Fitoterapia*, 2002; 73(7-8): 730–733
9. Judzientienė and J. Budienė, “Volatile constituents from aerial parts and roots of *Cichorium intybus* L. (chicory) grown in Lithuania,” *Chemija*, 2008; 19: 25–28.
10. Kim, S. B., I. S. Seo, M. A. Khan, K. S. Ki, M. S. Nam, and H. S. Kim. Separation of iron-binding protein from whey through enzymatic hydrolysis. *Int. Dairy J.*, 2007; 17: 625–631.
11. Kostias, G. Low-Fat and Delicious: Can we Break the Taste Barrier? *Journal of the American Dietetic Association*, 1997; 97: S88-S92.
12. D. Mares, C. Romagnoli, B. Tosi, E. Andreotti, G. Chillemi, and F. Poli, “Chicory extracts from *Cichorium intybus* L. as potential antifungals,” *Mycopathologia*, 2005; 160(1): 85–91.
13. Marie Celeste Karam<sup>1</sup>, Claire Gaiani<sup>1\*</sup>, Chadi Hosri<sup>2</sup>, Jennifer Burgain<sup>1</sup> and Joël Scher<sup>12013</sup>. Effect of dairy powders fortification on yogurt textural and sensorial properties: a review. *Journal of Dairy Research*, 2013; 80: 400–409.
14. Marshall, R., H. D. Goff, and R. W. Hartel. *Ice Cream*. Kluwer Academic Press, New York, NY, 2003.



15. M. C. Miller, S. K. Duckett, and J. G. Andrae, "The effect of forage species on performance and gastrointestinal nematode infection in lambs," *Small Ruminant Research*, 2011; 95(2-3): 188–192.
16. Nadia A. Abou-zeid Utilization of the Weed *Cichorium Pumilum*, Jacq as Vegetable Protease Enzymes for Whey Protein Hydrolysate Production *International Journal of Research Studies in Biosciences (IJRSB)*, 2015; 3(10): 44 – 53. ISSN 2349-0357 (Print) & ISSN 2349-0365 (Online).
17. S. Nandagopal and R. B. D. Kumari, "Phytochemical and antibacterial studies of chicory (*Cichorium intybus* L.)—a multipurpose medicinal plant," *Advances in Biological Research*, 2007; 1(1-2): 17–21.
18. Nielsen, B. J. Combined Emulsifier/Stabilizers for Ice Cream. *Gordian*, 1978; 78(6): 176-130.
19. Ohmes, R. L., Marshall, R. T. and O'Brien, C.M., et al. Evaluation of the effects of fat replacers on the quality of wheat bread, *Journal of Food Engineering*, 2003; 56: 265-267.
20. Phillips, L. G., J. B. German, T. E. O'Neill, E. A. Foegeding, V. Hanvalkar A., Kilara, B. A., Lewis, M. E., Mangino, C. V., Morr, J. M. Regenstein, D. M. Smith, and J. E. Kinsella. Standardized procedure for the measurement of the foaming properties of three proteins: A collaborative study. *J. Food Sci*, 1990; 55: 1441–1444-1453.
21. Plug, H., Haring, P. The role of ingredient-flavor interactions in the development of fat-free foods, *Trends in Food Science and Technology*, 1993; 4: 150-152.
22. P. Rani and N. Khullar, "Antimicrobial evaluation of some medicinal plants for their anti-enteric potential against multi-drug resistant *Salmonella typhi*," *Phytotherapy Research*, 2004; 18(8): 670–673.
23. C. Robert, T. H. Emaga, B. Wathelet, and M. Paquot, "Effect of variety and harvest date on pectin extracted from chicory roots (*Cichorium intybus* L.)," *Food Chemistry*, 2008; 108(3): 1008–1018.
24. M. A. Sanderson, M. Labreuveux, M. H. Hall, and G. F. Elwinger, "Nutritive value of chicory and English plantain forage," *Crop Science*, 2003; 43(5): 1797–1804.
25. Scharenberg, Y. Arrigo, A. Gutzwiller et al., "Palatability in sheep and in vitro nutritional value of dried and ensiled sainfoin (*Onobrychis viciifolia*) birdsfoot trefoil (*Lotus corniculatus*), and chicory (*Cichorium intybus*)," *Archives of Animal Nutrition*, 2007; 61(6): 481–496.
26. T. Shaikh, R.A. Rub, and S. Sasikumar, "Antimicrobial screening of *Cichorium intybus* seed extracts," *Arabian Journal of Chemistry*, 2012.



27. S'untar, E. K. Akkola, H.Kelesb, E.Yesiladac, S.D. Sarkerd, and T. Baykala, "Comparative evaluation of traditional prescriptions fromCichorium intybus L. forwound healing: stepwise isolation of an active component by in vivo bioassay and its mode of activity," Journal of Ethnopharmacology, 2012; 143(1): 299–309.
28. Q. Wang and J. Cui, "Perspectives and utilization technologies of chicory (Cichorium intybus L.): a review," African Journal of Biotechnology, 2011; 10(11): 1966–1977.