

REGULARIZATION BASED FUZZY GRAPH

Kailash Kumar Kakkad¹ and Dr. Sanjay Sharma*²

¹Department of Applied Mathematics, Chouksey Engineering College, Bilaspur (C. G.)
India.

²Department of Applied Mathematics, Bhilai Institute of Technology, Durg (C. G.) India.

Article Received on 24/01/2017

Article Revised on 14/02/2017

Article Accepted on 07/03/2017

***Corresponding Author**

Dr. Sanjay Sharma

Department of Applied
Mathematics, Bhilai
Institute of Technology,
Durg (C. G.) India.

ABSTRACT

The aim of this paper is introduce some definitions of regular fuzzy graphs, total degree and total regular fuzzy graph. Further we introduce concept of regularization based fuzzy graph and to obtain some result related to regular fuzzy graph and total regular fuzzy graph. Some basic theorems related to the stated graphs have also been presented.

AMS Subject Classification: 05C72.

KEYWORDS: Strong fuzzy graph, Regular fuzzy graphs, Totally regular fuzzy graph, Complete fuzzy graph, Degree of a vertex in fuzzy graphs.

INTRODUCTION

In 1736, the concept of graph theory was first introduced by Euler. The perception of fuzzy set was discussed by L.A. Zadeh, in 1965. In 1973, Kaufmann gave the first definition of a fuzzy graph which was based on Zadeh's fuzzy relations. A.Rosenfeld considered fuzzy relations on fuzzy sets and developed the theory of fuzzy graphs in 1975. Zadeh introduced the concept of fuzzy relations, in 1987. The theory of fuzzy graphs was developed J. N. Mordeson studied fuzzy line graphs and developed its basic properties, in 1993.

In this paper we discussed the concept of regular fuzzy graphs and totally regular fuzzy graphs. A comparative study between regular fuzzy graphs and totally regular fuzzy graph is made. Also some results on regular fuzzy graphs are studied and examined whether they hold for totally regular fuzzy graphs.

PRELIMINARIES

Definition 2.1: A graph $G^* = (V, E)$ is a pair of vertex set (V) and an edge set (E) where $E \subseteq V \times V$ i.e. E is a relation on V .

Definition 2.2: A fuzzy graph $G = (\sigma, \mu)$ is a pair of functions $\sigma : V \rightarrow [0, 1]$ and $\mu : V \times V \rightarrow [0, 1]$ with $\mu(x, y) \leq \sigma(x) \wedge \sigma(y)$, $\forall x, y \in V$, where V is a finite non- empty set and \wedge denote minimum. Where σ is a fuzzy subset of a non-empty set V and μ is a symmetric fuzzy relation on σ .

Definition 2.3: A fuzzy graph $G = (\sigma, \mu)$ is said to be a strong fuzzy graph if $\mu(x, y) = \sigma(x) \wedge \sigma(y)$, $\forall (x, y) \in E$.

Definition 2.4: A fuzzy graph $G = (\sigma, \mu)$ is said to be a complete fuzzy graph if $\mu(x, y) = \sigma(x) \wedge \sigma(y)$, $\forall x, y \in V$.


Definition 2.5: The degree of a vertex of G is denoted by

$$d_G(x) = \sum_{x \neq y} \mu(x, y)$$

REGULARIZATION ON FUZZY GRAPH

Definition 3.1: Let $G = (\sigma, \mu)$ be a fuzzy graph. If degree of each vertex is same say k , then G is called k -regular fuzzy graph.

Example 3.2: Let G be a fuzzy graph.



Here G is regular fuzzy graph, because degree of each vertex is $.8$, which is same.

Definition 3.3: Let $G = (\sigma, \mu)$ be a fuzzy graph. The total degree of a vertex of G is denoted by

$$Td_G(x) = \sum_{x \neq y} \mu(x, y) + \sigma(x) = d_G(x) + \sigma(x)$$

If total degree of each vertex is same say k , then G is called k - totally regular fuzzy graph.

Result 3.4

(I) Every regular fuzzy graph may or may not be a totally regular fuzzy graph.


(II) If fuzzy graph is not regular fuzzy graph, it may or may not be a totally regular fuzzy graph.

Result 3.5: If fuzzy graph is not regular, it may or may not be a totally regular fuzzy graph.

(i) If G is not regular fuzzy graph, G is also not totally regular fuzzy graph.

Example 3.6: Consider $G = (V, E)$ where $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and $E = \{v_1 v_2, v_2 v_3, v_3 v_4, v_4 v_5, v_5 v_6, v_6 v_1\}$

Define fuzzy graph G as below.



Then $d(v_1) = .4 + .2 = .6$, $d(v_2) = .2 + .5 = .7$, similarly $d(v_3) = .8$, $d(v_4) = .9$, $d(v_5) = .7$ and $d(v_6) = .5$. Since, $d(v_1) \neq d(v_2) \neq d(v_3) \neq d(v_4) \neq d(v_5) \neq d(v_6)$.

Therefore, G is not regular fuzzy graph.

But $Td(v_1) = .4 + .4 + .2 = 1.0$, $Td(v_2) = .2 + .1 + .5 = .8$, similarly $Td(v_3) = 1.5$, $Td(v_4) = 1.4$, $Td(v_5) = 1.3$, $Td(v_6) = .7$


Again $Td(v_1) \neq Td(v_2) \neq Td(v_3) \neq Td(v_4) \neq Td(v_5) \neq Td(v_6)$.

Therefore, G is not totally regular fuzzy graph.

(ii) G is not regular fuzzy graph but G is totally regular fuzzy graph.

Example 3.7: Consider $G = (V, E)$ where $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and $E = \{v_1 v_2, v_2 v_3, v_3 v_4, v_4 v_5, v_5 v_6, v_6 v_1\}$.

Define fuzzy graph G as below.



Then $d(v_1) = .3 + .2 = .5$, $d(v_2) = .2 + .4 = .6$, similarly $d(v_3) = .8$, $d(v_4) = .7$, $d(v_5) = .4$ and $d(v_6) = .4$. Since, $d(v_1) \neq d(v_2) \neq d(v_3) \neq d(v_4) \neq d(v_5) \neq d(v_6)$.

Therefore, G is not regular fuzzy graph. But $Td(v_i) = .9$ for all $i = 1, 2, 3, 4, 5, 6$.


Therefore, G is a totally regular fuzzy graph.

Result 3.8: Every regular fuzzy graph may or may not be a totally regular fuzzy graph.

(i) A fuzzy graph G is regular as well as totally regular fuzzy graph.

Example 3.9: Consider $G = (V, E)$ where $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and $E = \{v_1v_2, v_2v_3, v_3v_4, v_4v_5, v_5v_6, v_6v_1\}$.

Define fuzzy graph G as below.



Here $d(v_i) = .6 \forall i = 1, 2, 3, 4, 5, 6$


So G is regular. Also, $Td(v_i) = 1.1 \forall i = 1, 2, 3, 4, 5, 6$

$\therefore G$ is a totally regular fuzzy graph.

(ii) G is regular fuzzy graph, but it is not totally regular fuzzy graph.

Example 3.10: Consider $G = (V, E)$ where $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ and $E = \{v_1v_2, v_2v_3, v_3v_4, v_4v_5, v_5v_6, v_6v_1\}$.

Define fuzzy graph G as below.



Then $d(v_1) = 0.7 = d(v_2) = d(v_3) = d(v_4) = d(v_5) = d(v_6)$

So G is regular fuzzy graph.

But $Td(v_1) = .2 + .3 + .5 = 1.0$, $Td(v_2) = .5 + .5 + .2 = 1.2$, similarly $Td(v_3) = 1.1$, $Td(v_4) = .9$, $Td(v_5) = .8$, $Td(v_6) = .9$

So $Td(v_1) \neq Td(v_2) \neq Td(v_3) \neq Td(v_4) \neq Td(v_5) \neq Td(v_6)$.

$\therefore G$ is not totally regular fuzzy graph.

Theorem 3.11: Let σ be a constant function of fuzzy graph G . Prove that G is a regular fuzzy graph if G is a totally regular fuzzy graph.

Proof: Let σ be a constant function say c , i.e. $\sigma(x) = c$.

Let G be a k_1 – totally regular fuzzy graph.

$\therefore Td(x) = k_1$, for all $x \in V$.

Using definition of k_1 – totally regular fuzzy graph.

$\Rightarrow d(x) + \sigma(x) = k_1$, for all $x \in V$.

$\Rightarrow d(x) + c = k_1$, for all $x \in V$.

$\Rightarrow d(x) = k_1 - c$, for all $x \in V$.

$\therefore G$ is a regular fuzzy graph.

Thus G is a regular fuzzy graph if G is a totally regular fuzzy graph.

Theorem 3.12: Let σ be a constant function of fuzzy graph G . Prove that if G is a regular fuzzy graph, G is a totally regular fuzzy graph.

Proof: Let σ be a constant function say c , i.e. $\sigma(x) = c$.

Now, Let G be a k_2 – regular fuzzy graph.

Then $d(x) = k_2, \forall x \in V$.

Since we know that by definition of totally regular fuzzy graph.

$\therefore Td(x) = d(x) + \sigma(x), \forall x \in V$.

$\Rightarrow Td(x) = k_2 + c, \forall x \in V$.

Hence G is a totally regular fuzzy graph.

Thus if G is a regular fuzzy graph then G is a totally regular fuzzy graph.

Theorem 3.13: When a fuzzy graph G is regular as well as totally regular, then vertex set is a constant function.

Proof: Let σ be a vertex set and G be a k_1 –regular and k_2 – totally regular fuzzy graph.

So $d(x) = k_1$ and $Td(x) = k_2, \forall x \in V$.

Now $Td(x) = k_2, \forall x \in V$.

Using definition of k_2 – totally regular fuzzy graph.

$\Rightarrow \therefore d(x) + \sigma(x) = k_2, \forall x \in V$.

$\Rightarrow k_1 + \sigma(x) = k_2, \forall x \in V$.


$\Rightarrow \sigma(x) = k_2 - k_1, \forall x \in V$.

Hence σ is a constant function.

But converse of the above theorem 3.13 need not be true.

Example 3.15: Consider $G=(V,E)$ where $V= \{v_1,v_2,v_3,v_4,v_5,v_6\}$ and $E=\{v_1v_2,v_2v_3,v_3v_4,v_4v_5,v_5v_6,v_6v_1\}$.

Define fuzzy graph G as below



Then $d(v_1) = .7+.3 = 1.0$, $d(v_2) = .3+.4 = .7$ similarly $d(v_3) = .9$, $d(v_4) = .6$, $d(v_5) = .3$ and $d(v_6) = .9$

Since, $d(v_1) \neq d(v_2) \neq d(v_3) \neq d(v_4) \neq d(v_5) \neq d(v_6)$.

Therefore, G is not regular fuzzy graph.

But, $Td(v_1) = .7+.2+.3=1.2$, $Td(v_2) = .3+.4+.4=1.1$, similarly $Td(v_3) =1.4$, $Td(v_4) =1.1$, $Td(v_5) =.9$, $Td(v_6) =1.0$

Again $Td(v_1) \neq Td(v_2) \neq Td(v_3) \neq Td(v_4) \neq Td(v_5) \neq Td(v_6)$.

$\therefore G$ is not totally regular fuzzy graph.

Whereas σ is a constant function, because $\sigma(x) = Td(x) - d(x)$

$\sigma(v_1) = Td(v_1) - d(v_1)$

$\sigma(v_1) = 1.2 - 1.0$

$\sigma(v_1) = 0.4$, which is constant.

SUMMARY AND CONCLUSION

Therefore, there is no relation between regular fuzzy graphs and totally regular fuzzy graphs. A regular fuzzy graph may or may not be a totally regular fuzzy graph and in addition if fuzzy graph is not regular, it does not provide assurance for it to be a totally regular fuzzy graph.

REFERENCES

1. Vandana Bansal, "Effect of Regularization on Fuzzy Graph", 2016; IJAS, 017-020.
2. A. Nagoor Gani and S. R. Latha; "On Irregular fuzzy graphs" Applied Mathematical Sciences, 2012; 6(11): 517-523.
3. A. Nagoor Gani and K. Radha, "On Regular Fuzzy Graphs" Journal of Physical Sciences, 2010; 12: 33-40.
4. A. NagoorGani and K. Radha, Regular Property of Fuzzy Graphs, Bulletin of Pure and Applied Sciences, 2008; 27(2): 411-419.
5. L.A. Zadeh, Fuzzy Sets, Information and Control, 1965; 8: 338-353.
6. A. Rosenfeld, Fuzzy Graphs, In Fuzzy Sets and their Applications to Cognitive and Decision Process, Zadeh. L. A., Fu, K. S.,Shimura,M.,Eds; Academic Press, New York, 1975; 77-95.
7. Sunitha, M.S., and Vijayakumar, A.A.,Characterization of Fuzzy Trees, Information Sciences, 1999; 113: 293-300.

8. Bhutani, K.R., On Automorphism of fuzzy Graphs, Pattern Recognition Letters, 1991; 12: 413-420.
9. AL-Hawary, T., *Complete fuzzy graph*, International J.Math. Combin, 2011; 4, 26-34.
10. Johan N. Mordeson and Chang-Shyh Peng, operations on fuzzy graph, Information science, 1994; 79: 159-170.